

# STAT 510: Homework 04

David Dalpiaz

Due: Monday, March 1, 11:59 PM

## Exercise 1 (Make It So)

Let  $X_1, X_2, \dots, X_n \sim \text{Uniform}(0, \theta)$ . Consider the estimator

$$\hat{\theta} = \max\{X_1, X_2, \dots, X_n\}.$$

Find the bias, variance, and MSE of this estimator. Assuming the estimator is biased, create a new estimator which is a simple function of  $\hat{\theta}$  that is unbiased.

## Exercise 2 (More Data, Less Problems)

Let  $X_1, X_2, \dots, X_n \sim \text{Uniform}(0, \theta)$ . Consider the estimator

$$\hat{\theta} = 2 \cdot \bar{X}_n$$

Find the bias, variance, and MSE of this estimator. Is this estimator consistent? Justify.

## Exercise 3 (A Little Bit of Bias Goes a Long Way)

Let  $Y$  have a binomial distribution with parameters  $n$  and  $p$ . Consider two estimators for  $p$ :

$$\hat{p}_1 = \frac{Y}{n}$$

and

$$\hat{p}_2 = \frac{Y + 1}{n + 2}$$

For what values of  $p$  does  $\hat{p}_2$  achieve a lower mean square error than  $\hat{p}_1$ ?

## Exercise 4 (Dependence in the Empirical Distribution)

Let  $x$  and  $y$  be two distinct points. Find

$$\text{Cov}\left(\hat{F}_n(x), \hat{F}_n(y)\right).$$

### Exercise 5 (Empirical Distribution Properties)

For any fixed value of  $x$ , show each of the following.

$$\mathbb{E} \left[ \hat{F}_n(x) \right] = F(x)$$

$$\mathbb{V} \left[ \hat{F}_n(x) \right] = \frac{F(x) \cdot (1 - F(x))}{n}$$

$$\text{MSE} \left[ \hat{F}_n(x) \right] = \frac{F(x) \cdot (1 - F(x))}{n} \rightarrow 0$$

$$\hat{F}_n(x) \xrightarrow{P} F(x)$$

### Exercise 6 (Limiting Distribution of Empirical Distribution)

Let  $X_1, X_2, \dots, X_n \sim F$ . Given the empirical distribution function  $\hat{F}_n(x)$  and a fixed point  $x$ , use the central limit theorem to find the limiting distribution of  $\sqrt{n} \left( \hat{F}_n(x) - F(x) \right)$ .

### Exercise 7 (Using Statistical Functionals)

Let  $X_1, X_2, \dots, X_n \sim F$  and let  $\hat{F}_n(x)$  be the empirical distribution function. Let fixed numbers  $a < b$  and define

$$\theta = T(F) = F(b) - F(a).$$

Find the estimated standard deviation of

$$\hat{\theta} = T \left( \hat{F}_n(x) \right) = \hat{F}_n(b) - \hat{F}_n(a).$$

### Exercise 8 (More Coverage)

Let  $X_1, X_2, \dots, X_n \sim \text{Bernoulli}(p)$ . Set  $n = 100$  and  $\alpha = 0.05$ . Consider two confidence intervals for  $p$ . For both, define

$$\hat{p}_n = \frac{1}{n} \sum_{i=1}^n X_i.$$

First, consider the interval from the previous homework that we justified via Hoeffding's inequality.

$$C_n^H = \left( \hat{p}_n - \sqrt{\frac{1}{2n} \log \left( \frac{2}{\alpha} \right)}, \hat{p}_n + \sqrt{\frac{1}{2n} \log \left( \frac{2}{\alpha} \right)} \right)$$

Second, consider the "normal" interval,

$$C_n^N = \left( \hat{p}_n - z_{\alpha/2} \sqrt{\frac{\hat{p}_n(1 - \hat{p}_n)}{n}}, \hat{p}_n + z_{\alpha/2} \sqrt{\frac{\hat{p}_n(1 - \hat{p}_n)}{n}} \right).$$

Use simulation to check these intervals' coverage and expected length. Report your results using appropriate plots. Consider as many values of  $p$  as you can, but at minimum use

$$p \in (0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9).$$

Comment on the validity of these intervals and the interval lengths.

### Exercise 9 (Empirical Distribution Confidence Bands)

The following code simulates data from three different distributions.

```
set.seed(42)
data_1 = rexp(n = 100)
data_2 = rnorm(n = 25)
data_3 = rt(n = 500, df = 3)
```

For each, plot the empirical distribution with 95% confidence bands. For each, overlay the true cumulative distribution function. Do not use R's `ecdf()` function or anything similar. You may use R's `stepfun()` function.

### Exercise 10 (Estimating Functionals with the Empirical Distribution)

The following code simulates data from a [Weibull distribution](#).

```
set.seed(42)
some_data = rweibull(n = 250, shape = 2, scale = 3)
```

Use the empirical distribution function to create plug-in estimates of the following:

- Mean
- Variance
- Skewness
- Median

Compare these results to their true values given the data generating process defined above. Report your results as a table.

### Exercise 11 (Free Points)

Draw a smiley face!